

Original Research

Comparison Between Kinesiotape And Home-Based Exercise In Reducing Pain And Disability Due To Osteoarthritis In The Elderly

Putu Ayu Sita Saraswati^{1*}, Made Hendra Satria Nugraha²

^{1,2} Department of Physiotherapy, Faculty of Medicine, Udayana University, Indonesia

ABSTRACT

Background: OA in the knee causes various health problems, including pain and limited movement that reduces productivity and quality of life, causing disability. This study aims to determine the comparative effectiveness between home-based exercise and kinesiotape in lowering pain and disability in the elderly with knee osteoarthritis.

Methods: It was an experimental study with a randomized pre-test and post-test control group design. There were 30 participants divided into 2 groups: Group 1 with Kinesiotape and Group 2 with Home-Based Exercise. The intervention was given 3 times per week for 4 weeks. Disability was measured by WOMAC, and pain was measured by VAS.

Results: p-value for WOMAC and VAS scores before and after intervention is <0.001 in kinesiotape and Home-Based exercise interventions, which states that there is a significant improvement in lowering disability. Comparative analysis between groups also showed $p < 0.001$.

Conclusion: Kinesiotape and Home-Based Exercise are effective in lowering knee pain and disability in patients with knee pain due to genotype osteoarthritis. Home-based exercise is more effective than Kinesiotape in lowering knee pain and disability in patients with genotype osteoarthritis.

ARTICLE HISTORY

Received: November 12th, 2024

Accepted: May 5th, 2025

KEYWORDS

disability, exercise, kinesiotape, osteoarthritis, pain;

CONTACT

Putu Ayu Sita Saraswati



sitasaraswatui@unud.ac.id

Department of Physiotherapy,
Faculty of Medicine, Udayana
University. Jl. Bukit Jimbaran
Campus, Badung, Bali, Indonesia.

Cite this as: Saraswati, P. A. S., & Nugraha, M. H. S. (2024). Comparison Between Kinesiotape And Home-Based Exercise In Reducing Pain And Disability Due To Osteoarthritis In The Elderly. *Jurnal Keterapian Fisik*, 9(2), 1–12. <https://doi.org/10.37341/jkf.v9i2.454>

INTRODUCTION

Osteoarthritis (OA) is a degenerative disease caused by inflammation of the cartilage, bones, and synovium, making it the leading cause of disability in the elderly over 65 years old by 30% (Gao et al., 2023). Knee OA is one of the most common and debilitating forms of OA. It is often seen in the elderly, making it a significant health problem for this demographic.

The knee joint is a major weight-bearing joint, and over time, it can be subject to wear and tear, especially in older adults. As the cartilage in the knee deteriorates, it leads to inflammation, pain, and limited movement (Gelber, 2024). The factors that cause knee OA are gender, age, and occupation, which are related to the balance between bone degradation and hormones (Muhyi et al., 2023).

According to Data from the World Health Organization (WHO) in 2017, OA sufferers in the world approximately reached 9.6% in men and 18% in women (Brennan-Olsen et al., 2017). Basic Health Research (Riskesdas) in 2018 stated that the incidence of joint complaints in Indonesia was 7.3%, of which the majority of 8.5%, were reported by women. These complaints increase with increasing age, with a prevalence of 18.6% in people over 65 years old and 18.9% in people over 75 years old (Budiman & Widjaja, 2020).

Knee osteoarthritis (OA) in Indonesia affects 5% of individuals under 40 years old, 30% of those between 40 and 60 years old, and 65% of people over 61. Radiologically, the prevalence of knee OA is also notably high, with 15.5% of men and 12.7% of women showing signs of the condition. Global Burden of Disease (GBD) explained that from 1991-2019, there was a 114.5% increase in disability due to knee OA (Long et al., 2022). Osteoarthritis (OA) leads to a range of issues, including reduced physical function, changes in psychological well-being, restrictions in social interactions, challenges in fulfilling spiritual needs, and a decline in work productivity.

This has impact on the psychological, economic, and social status of OA complaints is very large, for the sufferer, his family, and the environment (Masyhurrosyidi et al., 2024). Pain is a physiological problem in the elderly with OA. The impact of pain on OA is fatigue, decreased range of motion, and pain when moving. There is also a severe morning stiffness when waking up. This causes a decrease in mobility, especially during extension, and limited physical mobility. It will result in disability and threaten the quality of life of the elderly (Masyhurrosyidi et al., 2024).

Physiotherapy treatment that can be provided for pain and disability conditions due to knee osteoarthritis in the elderly can be in the form of physiotherapy modality therapy and, exercise therapy, as well as the addition of kinesiotapes. Exercise therapy that can be done independently or home-based exercise and kinesiotape in standard physiotherapy interventions can have a positive effect in reducing pain and disability (Widnyana et al., 2024). Unfortunately, we found that home-based exercise for reducing the complaint needs further research, and less research has been conducted to prove its effect compared to kinesiotape.

Kinesiotape also has pro and cons impacts and still needs further research to prove its impact on osteoarthritis complaints. Those interventions need to be given to improve the quality of life of the elderly. However, this type of home-based exercise is rarely applied. Researchers are interested in conducting further studies on this matter. This study aims to determine the comparative effectiveness between home-based exercise and kinesiotape in lowering pain and disability in the elderly with knee osteoarthritis.

MATERIALS AND METHODS

This study is an experimental design featuring a randomized pre-test and post-test control group. Its goal is to compare the effects of kinesiotaping and home-based exercise on reducing disability caused by knee osteoarthritis (OA). The research took place in private physiotherapy clinics in Denpasar and Badung, Bali, from July to September 2024. Participants were randomly assigned using the permutation block method.

Sampling was conducted according to the inclusion and exclusion criteria. Inclusion criteria include individual 60 years old and above; having knee pain almost every day in the past month due to knee OA according to a doctor's referral; average pain score of 3 and 7 on NRS 10 points; have good cognitive function MMSE Score 24-

30; and able to walk independently on flat surfaces without assistive devices. The exclusion criteria for this study include individuals with a history of joint replacement or knee arthroscopic surgery, those who have had other surgeries on the lower extremities within the past six months, and individuals with severe deformities of the lower extremities (such as varus or valgus knee).

Additionally, participants with health conditions that could lead to side effects during home exercise, including uncontrolled high blood pressure, myocardial infarction, cerebral infarction, unstable angina, arrhythmias, severe vision issues, or neurological dysfunction, are also excluded. The withdrawal criteria consist of: (a) the patient failing to return during the study, (b) a deterioration in the patient's condition after receiving therapy, and (c) voluntary resignation from the study. Sampling was conducted through a physiotherapy assessment, along with a doctor's referral indicating that the patient experienced pain and disability due to knee osteoarthritis.

The sample size is used by the G*Power application. The effect size was determined through research (Yilmaz et al., 2019). The α value of error probability is set at 0.05, and power ($1 - \beta$ error probability) is set at 0.80. Based on the calculation results, the number of subjects was 13. To avoid dropping out, subjects were added by 20% to 14.3 or 15 research participants in each group. The total number of research participants in both groups is 30.

Kinesiotape and home-based exercise were the independent variables, while the dependent variables were pain and disability associated with osteoarthritis (OA). The research procedure to achieve the research objectives is as follows: the researcher conducts a licensing process at the institution where the research is conducted with the number 2191/UN14.2.2.VII.14/LT/2024. The researcher prepares an informed consent form, which must be signed by the participant and approved by the physiotherapy supervisor.

This form confirms the participant's willingness to take part in the study until its completion. Additionally, the researcher provides educational information to the participants being researched about the benefits, objectives, how this research is conducted, and the importance of conducting this research. The application technique of kinesiotape used is in line with a study from Donec & Kubilius (2019), which uses two Y-shapes and two I-shapes. Home-based exercise involves 10 types of lower limb strengthening exercises.

The measuring tools evaluated in this study were pain and disability. Pain was measured by the Visual Analogue Scale, which had a good to excellent validity score on the relationship between VAS and NRS ($r=0.941$) and VAS and VRS ($r=0.878$), while the reliability score had an excellent value ($r=0.97$) on the test-retest reliability (Alghadir et al., 2018). Knee disability was measured using the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) in the Indonesian Version, which had a good validity based on the assessment of orthopaedics experts and a high internal consistency score, tested with Cronbach's Alpha, is 0.966 (Karsten et al., 2019).

After 6 weeks of exercise intervention and 12 intervention sessions and the researcher obtained complete data, then the researcher compared the results before and after the intervention in both groups with treatment and conducted a differential test. In analyzing the data obtained from disability measurements, changes in the level of pain reduction before and after the intervention were analyzed using the SPSS (Statistical Program for Social Science) program. The normality of WOMAC and Pain (VAS) scores was assessed using the Shapiro-Wilk test. The p-values for both scores in both

groups were greater than 0.05, indicating that the data in both groups followed a normal distribution both before and after treatment.

RESULTS

The characteristics of the research subjects include age, gender, height, and weight. The description of the characteristics of the research subjects is presented in Table 1.

Table 1. Subjects Characteristics

Characteristics	Group 1	Group 2
Age (years)	65.13±3.4	65.4 ±3.1
Gender		
Male	5 (33.3)	7 (46.7)
Female	10 (66.7)	8 (53.3)
Height (Cm)	161.5 ±6.9	159.9±5.9
Weight (Kg)	63.3±9.5	66.5±11.1

Based on Table 1 shows that the research subjects in Group 1 have an average age of 65.13 years, and Group 2 has an average age of 65.4 years. In group 1, there were 10 female subjects (66.7%), and more than 5 men (33.3%). In group 2, the subjects were male as many as 7 people (46.7%) and female as many as 8 people (53.3%). It can be concluded that the sample with female sex is more than the male sample in both group 1 and group 2. Based on Table 1 it shows that the research subjects in group 1 have an average height of 161.5 cm with an average weight of 63.3 kg, while group 2 with an average height of 159.9 cm and a body weight of 66.5 kg.

Table 1 shows that the number of knee OA sufferers is higher in women than in men. By the findings of this study, the prevalence of knee pain due to OA in Indonesia is experienced in the elderly population, where women have a higher percentage of 8.5%, while men have 6.1% of the total percentage of 17.3% (Muhyi et al., 2023). The cause of the high percentage experienced by elderly women is most often related to hormonal influences (Soeryadi et al., 2017). Elderly women will experience a menopause phase, which is an inactive reproductive organ which is marked by the cessation of the menstrual cycle.

The cessation of this menstrual cycle causes a decrease in various hormones, especially the hormone estrogen. The hormone estrogen acts as an inhibitor of bone resorption by osteoclasts (Sihombing et al., 2013). The decrease in the hormone estrogen in elderly women results in the task of osteoclasts that continue to be active cannot be stopped, accelerating calcification in the joints. The difference with male elderly who are still active produces testosterone hormone, which functions in inhibiting osteoclasts, so that female elderly are susceptible to OA in the knee if monitored in gender difference studies (Suari et al., 2015).

Weight is one of the factors that cause knee OA. Elderly women experience obesity faster after menopause, due to the hormone estrogen, which decreases its function in reducing body fat deposits (Soeryadi et al., 2017). The occurrence of pressure by excess weight on the knee joint accelerates the cartilage damage, which will cause friction and a crepitating sound if there is movement of the knee joint when walking (Suari et al., 2015). In this study, weight and height measurements were carried out to be able to evaluate the relationship between these factors in knee OA.

Table 2. Results of the Homogeneity Test of WOMAC Score and Pain in OA Genu Before and After Treatment

Data Group	Homogeneity Test
WOMAC Before Intervention	0.554
WOMAC After Intervention	0.871
VAS Before Intervention	0.719
VAS After Intervention	0.359

Based on Table 2, the results of the homogeneity test using Levene's test obtained a probability value for the WOMAC score value of the data group before treatment, which is $p = 0.554$, and the data value after treatment, which is $p = 0.871$. The results showed that the variation of WOMAC scores before and after treatment between the two groups was homogeneous ($p > 0.05$). In the pain value based on VAS, the result before treatment is $p = 0.719$, and after treatment, which is $p = 0.359$, showing a p value > 0.05 , which means that the value of VAS data before and after treatment is homogeneous.

To determine the decrease in disability before and after the intervention in the group, a two-sample mean difference test was carried out from the two datasets with a paired sample t-test. The test results are shown in Table 3.

Table 3. Effect of Intervention on Reduction of Disability and Pain in OA Genu in the Elderly in Each Group

Data Group	Before Intervention Mean \pm SD	After Intervention Mean \pm SD	p value
VAS			
Group 1	5,35 \pm 0,49	2,82 \pm 0,96	<0,001
Group 2	5,29 \pm 0,56	2,03 \pm 0,46	<0,001
WOMAC			
Group 1	30,20 \pm 5,1	19 \pm 3,66	<0,001
Group 2	28,73 \pm 4,78	13 \pm 3,30	<0,001

Table 3 shows the results of the hypothesis test conducted using the Independent T-test to determine the difference in knee disability in group 1. The calculation results obtained p-values for the WOMAC and VAS scores are <0.001 , which states that disability and knee pain decrease significantly before and after the Kinesiotape intervention ($p < 0.001$). Hypothesis testing to determine the difference in knee disability before and after treatment in group 2 also uses the Independent T-Test difference test. The calculation results obtained p-values for the WOMAC and VAS scores are <0.001 , which states that there is a significant difference in the reduction of disability and knee pain because of the Home-Based exercise intervention.

Table 4. Effect of Intervention on Reduction of Disability and Pain in OA Genu in the Elderly Between Group

	Data Group	Mean \pm SD	p value
Pre-Test	Group 1	5,35 \pm 0,49	0,759
(VAS)	Group 2	5,29 \pm 0,56	
Post-Test	Group 1	2,82 \pm 0,96	<0,001
(VAS)	Group 2	2,03 \pm 0,46	
Difference	Group 1	2,52 \pm 0,79	0,424

	Data Group	Mean \pm SD	p value
(VAS)	Group 2	3,26 \pm 0,34	
Pre-Test	Group 1	30,20 \pm 5,1	
(WOMAC)	Group 2	28,73 \pm 4,78	
Post-Test	Group 1	19 \pm 3,66	
(WOMAC)	Group 2	13 \pm 3,30	<0,001
Difference	Group 1	11,20 \pm 2,5	
(WOMAC)	Group 2	15,73 \pm 3,2	

The results of Table 4 show that the p-value of the difference between the two variables after treatment between groups is <0.001. These results indicate that there is a significant difference in the results of the implementation of the intervention in the two groups (p<0.001). From the table results, it can also be concluded that Home-Based Exercise reduces knee disability more than Kinesiotape, this can be seen in the results of group 2, which are closer to the p-value.

DISCUSSION

Reduction of Knee Disability and Pain due to OA Genu in the Elderly with Kinesiotape Intervention

From the data test of the results of group 1 research, there was a significant difference in the results before and after the kinesiotape intervention on the WOMAC and VAS scores of the subjects. Kinesiotape is a rehabilitative therapy using an elastic tape made of polymer that can be stretched 120-140% (Azizah et al., 2024). The thickness and elasticity of the kinesiotape are adjusted like human skin, so it will not limit the scope of motion or put excessive emphasis on the adhesive area (Yuri et al., 2022).

The kinesiotape method was invented by Dr. Kase from Japan in the 1970s. The application of kinesiotape is in direct contact with the skin and left on for several hours with good adhesion and low risk of skin irritation. The good or bad results of kinesiotape depend on the application techniques used, such as Y, I, and Fan Shaped (Donec & Kubilius, 2019).

The application technique used is in line with a study from Donec & Kubilius (2019), which uses two Y-shapes and two I-shapes. Two of the Y-shaped techniques are applied to the vasodilation of blood vessels and lymphatics in overcoming effusion of the knee joint. Following this, an I-shaped strip is placed just below the lower boundary of the patella, above the patellar tendon, while the knee is fully bent and the subject is in a supine position, applying 100% tensile tension along with adhesive activation.

The knee is then positioned at a 20–30° bend, and the adhesive is continued over the medial and lateral collateral ligaments, using about 75% tensile tension with adhesive activation. After this, the subject is instructed to fully straighten the knee, with the tip of the I-shaped strip (approximately 10 cm) directed to the posterolateral side of the thigh (ensuring no overlap at the back) with 0% tension and adhesive activation. A second I-shaped strip is then applied in the same manner, ensuring it does not overlap the first one.

According to the study by Donec and Kubilius (2019), there was a significant reduction in generalized knee pain, including pain at night, during the day, when changing body positions, and during prolonged activities such as running, climbing stairs, and walking (p < 0.05). At rest, the pain reduction was similar for participants in

both groups ($p = 0.421$). Kinesiotape was applied for three days, with periodic replacements done three times a week.

This protocol implies that kinesiotaping is intended to be used regularly to maintain its effects. The periodic replacement of the tape ensures that it remains effective in providing support and alleviating pain. Given the regularity of tape replacement, it can be inferred that continuous and consistent use is necessary for maintaining pain relief and potentially improving the long-term outcomes for individuals with knee pain.

In a study conducted by Pamuk and Yucesoy (2015), Magnetic Resonance Imaging (MRI) was used to assess issues in the limb after kinesiotape was applied over the skin along the tibialis anterior muscle with a pull of approximately 50%. The study indicated that kinesiotape exerts mechanical effects not only on the superficial tissues but also on deeper layers, leading to heterogeneous deformation throughout the limb. Ruffini corpuscles and interstitial receptors are found in the boundary zone of soft tissue layers.

Applying kinesiotape with a slight pull in a specific direction can enhance afferent stimulation and decrease skin friction. Additionally, a study by Castrogiovanni et al. (2016) reported that the application of kinesiotape with tensile tension resulted in a more significant long-term analgesic effect compared to application without tension. The Visual Analog Scale (VAS) pain scores indicated that, across all groups, patients exhibited lower scores after both 15 days and 3 months ($p < 0.05$). However, in Group 3, the scores remained unchanged between the 15-day and 3-month marks ($p = \text{n.s.}$).

The kinesiotape intervention given to group 1 with knee OA functioned to provide stability to the joints with a micro-lifting mechanism to help increase blood and lymphatic circulation, vasodilation, and reduce the occurrence of pressure by body weight on inflamed joint structures (Yuri et al., 2022). The occurrence of vasodilation of blood vessels creates a metabolic system in the knee, causing inflammatory substances such as bradykinin, prostaglandins, and histamine to be dissolved and washed with the bloodstream, so that local pain will decrease. Stabilization of the installed kinesiotape provides a proprioceptive stimulus of sensory feedback to help the subject be more aware of the position of the knee joint, which can later reduce the risk of unwanted movements or worsen disability (Azizah et al., 2024).

This study is in line with (Widnyana et al., 2024), namely the addition of kinesiotape after intervention modalities of ultrasound therapy, transcutaneous electrical nerve stimulation, and massage, proven to be effective in improving functional ability in patients with knee OA. In this study, an independent t-test was used, which stated that there was a significant difference in WOMAC scores between the two groups before and after the intervention, with a value of $p < 0.05$. Functional ability in the treatment group increased by 30.57% from the conclusion of kinesiotape addition after the intervention.

The mechanism of action of kinesiotape in improving functional ability here with sensory responses to the skin in the area of the muscle belly and attached tendons can increase the strength of the quadriceps muscles. The concentric contraction generated by the kinesiotape pull is fitted with a facilitation technique from origo to insorcio, with a 40% pull will recruit more unit motors. We know that in the elderly, changes in the posture of the "X" and "O" shaped legs often occur; the function of the kinesiotape as a stabilizer can maintain the position of the patella according to its structure, so that disability in the knee can be reduced.

Reduction of Knee Disability and Pain due to OA Genu in the Elderly with Home-Based Exercise Intervention

In the current study, the Home-Based Exercise (HBE) intervention given in group 2 showed a significant difference in the reduction of knee disability before and after the intervention ($p < 0.001$). HBE is a collection of several trainings that can be done independently at home to reduce pain, increase muscle strength, flexibility, and maintain the scope of motion of the knee joint (Si et al., 2023). Without the need for expensive tools or large costs, HBE can be done efficiently and only requires leisure time and a convenient place to do it (Mao et al., 2024).

Knee OA sufferers are also characterized by the presence of thigh muscle weakness of up to 20-45%, depending on age. In the knee joint, the quadriceps muscles as an important protective function, that is working in eccentric conditions during the initial standing phase of walking to dampen the knee joint and acting to slow down the limb before the heel hits the ground, thus reducing excessive impulsive loads. Weakness in the quadriceps muscle can be attributed to increased load on the knee joint, loss of patellofemoral cartilage, and narrowing of the tibiofemoral joint space.

According to a study by Rice et al., (2011) that examined the mechanism of quadriceps muscle weakness in knee OA, this weakness is Arthrogenic Muscle Inhibition (AMI), an ongoing nerve block that prevents the quadriceps muscle from being fully activated. AMI can cause muscle atrophy in the quadriceps. A characteristic feature seen in knee OA is stiffness in the joints. Joint stiffness can be described as discomfort or difficulty during movement because the joint is perceived as inflexible and restricted. A decrease in the amount of Surface-Active Phospholipid (SAPL) production in the synovial membrane results in the erosion of cartilage in the joint.

As a result, friction occurs between the bones of the joints and inhibits the movement of the joint's range of motion (Widnyana et al., 2024). Early pathological changes in knee osteoarthritis (OA) primarily occur on the surfaces of articular cartilage in areas experiencing the highest loading stress. In response to matrix loss, there is an increase in chondrocyte proliferation. Some chondrocytes transform into hypertrophic chondrocytes, resembling those found in the hypertrophic zone of the growth plate.

During the degradation phase of OA, substantial matrix loss takes place due to ongoing protease production, which is stimulated by proinflammatory cytokines. These cytokines encourage chondrocytes to generate even more cytokines and proteases. Furthermore, significant matrix damage results from cell loss, largely due to chondrocyte apoptosis (Heika et al., 2020).

Many of the issues related to knee osteoarthritis (OA) have been addressed in this study, which shares objectives with the research by Si et al., (2023). The primary goals include enhancing thigh muscle strength, increasing the knee joint's range of motion, and improving muscle and joint flexibility. The program is implemented in stages, beginning with an initial training phase that involves four sessions of approximately 30 minutes each over the first two weeks, followed by follow-up appointments at monthly intervals. Participants are encouraged to perform the exercises for both legs for 20-30 minutes daily, gradually increasing to a maximum of 20 repetitions per leg. Adherence to the program is monitored through a self-reported diary, collected weekly.

The exercises prescribed include: (1) chair sitting; (2) isometric hamstring exercises; (3) isotonic hip extension exercises while lying down; (4) isometric hip adductor exercises while sitting; (5) isotonic hip abductor exercises while lying down; (6) side-lying isotonic hip abductor exercises; (7) narrow stance squats; (8) wide stance

squats; (9) hamstring stretches while sitting; and (10) quadriceps stretches while lying on the side.

Research by Mao et al. (2024) found similar results, indicating that HBE significantly reduces pain (SMD=-0.38, 95% CI (-0.58, -0.18); P = 0.001), enhances joint function (SMD=-0.60, 95% CI (-1.01, -0.19); P = 0.004), improves balance (SMD=-0.67, 95% CI (-1.00, -0.34); P = 0.001), and increases mobility in activities of daily living (ADL) (SMD=0.51, 95% CI (0.19, 0.82); P = 0.002) for patients with knee OA. Thus, it can be concluded that HBE not only alleviates complaints associated with knee OA but also enhances walking quality. However, its effectiveness depends on consistent and regular practice (Mao et al., 2024).

Comparison of Kinesiotape and Home-Based Exercise in Reducing Knee Disability and Pain due to OA Genu in the Elderly

Based on the research that has been done, the administration of Home-Based Exercise is more effective compared to kinesiotape. HBE has the advantage of being easy to apply independently, improving disability function, reducing pain, and having no need to spend money (Castrogiovanni et al., 2016). However, HBE has shortcomings, namely the need for consistency and compliance in undergoing training and the need for supervision and assistance from the family, where the subject here is an elderly person. In addition, the ability to be flexible and the strength of the elderly will certainly affect the intervention of HBE (Castrogiovanni et al., 2016).

Kinesiotape interventions have the advantage of accelerating the reduction of joint inflammation, reducing pain, and stabilizing the knee joint. However, the application of kinesiotape requires consumables, and installation must also be monitored to be accurate and correct. The study did not include a long-term follow-up period to assess the sustainability of pain relief after the kinesiotaping was discontinued. It would be valuable to evaluate whether the benefits of kinesiotaping persist over time or if pain levels return once treatment stops.

CONCLUSION

Based on the analysis of research that has been carried out and the discussion, it can be concluded that home-based exercise and kinesiotape are effective in reducing knee pain and disability in patients with knee pain due to osteoarthritis. Home-based exercise is found to be more effective than kinesiotape in reducing knee pain and disability in patients with knee pain due to genotype osteoarthritis. ... The intervention period was limited to three days, and the effects of kinesiotaping on knee pain over the long term were not evaluated. Future studies should explore whether the pain relief observed in the short term can be maintained over extended periods of use.

ACKNOWLEDGEMENT

The author would like to thank the Institute for Research and Community Service, Udayana University, and the Research and Community Service Unit, Faculty of Medicine, Udayana University, for the 2024 Study Program Leading Research Grant (Hibah Penelitian Unggulan Program Studi tahun 2024).

REFERENCES

Alghadir, A., Anwer, S., Iqbal, A., & Iqbal, Z. (2018). Test-retest reliability, validity, and minimum detectable change of visual analog, numerical rating, and verbal

rating scales for measurement of osteoarthritic knee pain. *Journal of Pain Research*, Volume 11, 851–856. <https://doi.org/10.2147/JPR.S158847>

- Azizah, N., Sam, D., & Kuswardhana. H. (2024). Pengaruh Terapi Kinesio Taping terhadap Penurunan Nyeri Penderita Osteoarthritis . *Jurnal Pendidikan Tambusai*, 8(1), 11058–11068.
- Brennan-Olsen, S. L., Cook, S., Leech, M. T., Bowe, S. J., Kowal, P., Naidoo, N., Ackerman, I. N., Page, R. S., Hosking, S. M., Pasco, J. A., & Mohebbi, M. (2017). Prevalence of arthritis according to age, sex and socioeconomic status in six low and middle income countries: analysis of data from the World Health Organization study on global AGEing and adult health (SAGE) Wave 1. *BMC Musculoskeletal Disorders*, 18(1), 271. <https://doi.org/10.1186/s12891-017-1624-z>
- Budiman, N. T., & Widjaja, I. F. (2020). Gambaran derajat nyeri pada pasien osteoarthritis genu di Rumah Sakit Royal Taruma Jakarta Barat. *Tarumanagara Medical Journal*, 2(2), 372–377. <https://doi.org/10.24912/tmj.v3i1.9744>
- Castrogiovanni, P., Di Giunta, A., Guglielmino, C., Roggio, F., Romeo, D., Fidone, F., Imbesi, R., Loreto, C., Castorina, S., & Musumeci, G. (2016). The Effects of Exercise and Kinesio Tape on Physical Limitations in Patients with Knee Osteoarthritis. *Journal of Functional Morphology and Kinesiology*, 1(4), 355–368. <https://doi.org/10.3390/jfmk1040355>
- Donec, V., & Kubilius, R. (2019). The effectiveness of Kinesio Taping® for pain management in knee osteoarthritis: a randomized, double-blind, controlled clinical trial. *Therapeutic Advances in Musculoskeletal Disease*, 11. <https://doi.org/10.1177/1759720X19869135>
- Gao, B., Li, L., Shen, P., Zhou, Z., Xu, P., Sun, W., Zhang, C., & Song, Q. (2023). Effects of proprioceptive neuromuscular facilitation stretching in relieving pain and balancing knee loading during stepping over obstacles among older adults with knee osteoarthritis: A randomized controlled trial. *PLOS ONE*, 18(2), e0280941. <https://doi.org/10.1371/journal.pone.0280941>
- Gelber, A. C. (2024). Knee Osteoarthritis. *Annals of Internal Medicine*, 177(9), ITC129–ITC144. <https://doi.org/10.7326/ANNALS-24-01249>
- Karsten, S., Limena, S., & Phandu, M. (2019). Translation, adaptation, and validation of western ontario and mcmaster universities osteoarthritis index (WOMAC) for indonesian. *Jurnal Orthopaedi Dan Traumatologi Indonesia*, Volume 2 Issue 3. <https://doi.org/10.31282/joti.v2n3.48>
- Long, H., Liu, Q., Yin, H., Wang, K., Diao, N., Zhang, Y., Lin, J., & Guo, A. (2022). Prevalence Trends of Site-Specific Osteoarthritis From 1990 to 2019: Findings From the Global Burden of Disease Study 2019. *Arthritis & Rheumatology*, 74(7), 1172–1183. <https://doi.org/10.1002/art.42089>

- Mao, Y., Qiu, B., Wang, W., Zhou, P., & Ou, Z. (2024). Efficacy of home-based exercise in the treatment of pain and disability at the hip and knee in patients with osteoarthritis: a systematic review and meta-analysis. *BMC Musculoskeletal Disorders*, 25(1), 499. <https://doi.org/10.1186/s12891-024-07585-w>
- Masyhurrosyidi, H., Kumboyono, & Utami, Y. (2024). Pengaruh Kompres Hangat Rebusan Jahe Terhadap Tingkat Nyeri Subakut dan Kronis pada Lanjut Usia dengan Osteoarthritis Lutut di Puskesmas Arjuna Kecamatan Klojen Malang Jawa Timur. *Majalah Kesehatan*, 1(1), 39–44.
- Muhyi, A., Adiratna, B., & Pertiwi, S. (2023). Prevalensi Osteoarthritis Genu Berdasarkan Karakteristik Demografi pada Pasien Geriatri di RSUD K.R.M.T. Wongsonegoro. *Jurnal Kesehatan Masyarakat*, 11(2), 152–160.
- Rice, D. A., McNair, P. J., & Lewis, G. N. (2011). Mechanisms of quadriceps muscle weakness in knee joint osteoarthritis: the effects of prolonged vibration on torque and muscle activation in osteoarthritic and healthy control subjects. *Arthritis Research & Therapy*, 13(5), R151. <https://doi.org/10.1186/ar3467>
- Si, J., Sun, L., Li, Z., Zhu, W., Yin, W., & Peng, L. (2023). Effectiveness of home-based exercise interventions on pain, physical function and quality of life in individuals with knee osteoarthritis: a systematic review and meta-analysis. *Journal of Orthopaedic Surgery and Research*, 18(1), 503. <https://doi.org/10.1186/s13018-023-04004-z>
- Sihombing, I., Wangko, S., & Kalangi, S. J. R. (2013). Peran Estrogen pada Remodeling Tulang. *Jurnal Biomedik (JBM)*, 4(3). <https://doi.org/10.35790/jbm.4.3.2012.1210>
- Soeryadi, A., Gesal, J., & Sengkey, L. S. (2017). Gambaran Faktor Risiko Penderita Osteoarthritis Lutut di Instalasi Rehabilitasi Medik RSUP Prof. Dr. R. D. Kandou Manado Periode Januari –Juni 2017. *E-CliniC*, 5(2). <https://doi.org/10.35790/eci.5.2.2017.18540>
- Suari, B., Ihsan, M., & Burhanuddin, L. (2015). *Gambaran Penderita Osteoarthritis di Bagian Bedah RSUD Arifin Achmad Periode Januari 2011 - Desember 2013*. <https://Media.Neliti.Com/Media/Publications/183923-ID-Gambaran-Penderita-Osteoarthritis-Di-Bagi.Pdf>.
- Widnyana, M., Pramana Putra, I. P. Y., Septian Utama, A. A. G. E., & Kamayoga, I. D. G. A. (2024). Penambahan Kinesiotaping pada Intervensi Ultrasound, Transcutaneous Electrical Nerve Stimulation, dan Massage lebih Meningkatkan Kemampuan Fungsional pada Pasien Osteoarthritis Genu. *Jurnal Fisioterapi Dan Rehabilitasi*, 8(2), 234–241. <https://doi.org/10.33660/jfrwhs.v8i2.379>
- Yilmaz, M., Sahin, M., & Algun, Z. C. (2019). Comparison of effectiveness of the home exercise program and the home exercise program taught by physiotherapist

in knee osteoarthritis. *Journal of Back and Musculoskeletal Rehabilitation*, 32(1), 161–169. <https://doi.org/10.3233/BMR-181234>

Yuri, S. A., Hadi, P., & Faridah, F. (2022). Penatalaksanaan Fisioterapi pada Kasus Osteoarthritis Genu Dextra dengan Metode Kinesiotapping, Hold Relax Stretching, dan Modalitas US. *Jurnal Kajian Ilmiah Kesehatan Dan Teknologi*, 4(1), 1–7. <https://doi.org/10.52674/jkikt.v4i1.55>